

# Regeneration of Ammonia Borane Spent Fuel(s) by Direct Reaction with Hydrazine and Liquid Ammonia

Andrew D. Sutton, Anthony K. Burrell, David A. Dixon,  
Edward B. Garner III, John C. Gordon, Tessui Nakagawa,  
Kevin C. Ott, J. Pierce Robinson, Monica Vasiliu.




Operated by Los Alamos National Security, LLC for the U.S. Department of Energy's NNSA



# What Does H<sub>2</sub> Offer as an Alternative Fuel?

- Potentially part of a renewable energy cycle
- Water is the sole oxidation product



- Favorable energy content per kg (120 MJ) compared to petroleum (44 MJ)


(corresponds to ~19 MJ/L for AB (Based on 3H<sub>2</sub> released) versus ~34MJ/L for gasoline)

- Can be used in a PEM fuel cell

# How Can H<sub>2</sub> be Stored in a Vehicle?

- Physical
  - As a compressed gas at high pressure
  - As a liquid, cryogenically
  - Using sorbants (nanoporous materials)
- Chemical
  - Metal hydrides
  - Chemical hydrides

# DOE Volumetric and Gravimetric Targets<sup>1</sup>




Relative volumes of 4 kg of H<sub>2</sub><sup>2</sup>

|                     | 2010 | 2015 | Ultimate |
|---------------------|------|------|----------|
| Gravimetric (wt. %) | 4.5  | 5.5  | 7.5      |
| Volumetric (MJ/L)   | 3.2  | 4.7  | 8.3      |

<sup>1</sup>[http://www1.eere.energy.gov/hydrogenandfuelcells/storage/pdfs/targets\\_onboard\\_hydro\\_storage.pdf](http://www1.eere.energy.gov/hydrogenandfuelcells/storage/pdfs/targets_onboard_hydro_storage.pdf)

<sup>2</sup>Schlapbach, L.; Züttel, A. *Nature* **2001**, 414, 353-358.


# Why Ammonia Borane?



Ammonia Borane (AB)

- Advantages
  - Non-toxic
  - Hydridic and protic hydrogen atoms
  - High wt. % of hydrogen (6.5/13.1/19.6)
- Challenges
  - Controlled Dehydrogenation
  - Regeneration



# Methods of Dehydrogenation





# Earth Abundant Catalyst Development



Colorless solid, ca. 18 % purified yield  
Cy = Cyclohexyl, TMS = Trimethylsilyl




$\text{PCy}_3\text{Fe}[\text{N}(\text{TMS})_2]_2 + \text{AB}$  in THF at RT.  
P6<sub>3</sub>, R1 = 11.5 %, 50 % thermal ellipsoids, cyclohexyl P substituents omitted.



- Mixture of products
  - Depends on rxn. conditions
- 2.5 – 3 h reaction time
- ca. 3.5 mol. % catalyst loading
- 1.5 – 1.7 eq. H<sub>2</sub> per AB
- Very sensitive to oxidation/decomp.
- Trigonal planar coordination geom.

# Off-Board Regeneration Required



$$\Delta H \approx -7 \text{ kcal/mol}$$

(Miranda and Ceder 2007)

Hydrogen release -> **too exothermic** to  
merely re-pressurize with H<sub>2</sub>




Off-board chemistry required to  
regenerate ammonia borane (AB)

N.B. Within the Center we are working on systems that are potentially  
directly regenerable (on-board??):


e.g. Ca(NH<sub>2</sub>BH<sub>3</sub>)<sub>2</sub> (A.K. Burrell et. al. *Angew. Chem. Int. Ed.* **2007**, 46, 8995).






# Regeneration Strategy



# Benzenedithiol Chemistry Works with Borazine and PB...



# Sn-H Recycle from Sn-S is Also Required...



# Drive to Reduce Mass - 100 Metric Ton/Day Recycle of AB.....


## Rohm & Haas Report Detail

|   | Digestion      | Metal Reduction<br>/ Amine<br>Exchange | Ammoniation    | Metal Recovery   |
|---|----------------|--|----------------|------------------|
| H <sub>2</sub>  |                | 431                                    |                | 1,145            |
| NH <sub>3</sub>   |                | 15,645                                 | 3,303          |                  |
| (Et) <sub>2</sub> NH  |                | 14,930                                 | 67,187         |                  |
| THF   | 86,989         | 86,989                                 |                |                  |
| Toluene   |                |  | 88,773         |                  |
| BNH   | 5,272          |  |                |                  |
| C <sub>6</sub> H <sub>4</sub> (SH) <sub>2</sub>                                 | 22,505         | 2,250                                  |                | 222,795          |
| Bu <sub>3</sub> SnH   |                | 162,797                                |                | 906,777          |
| (Et) <sub>2</sub> NHBH <sub>3</sub>   |                | 79,896                                 | 8,877          |                  |
| HB(C <sub>6</sub> H <sub>4</sub> S <sub>2</sub> )NH <sub>3</sub>                | 69,895         | 17,258                                 |                | 17,258           |
| (NH <sub>4</sub> )B(C <sub>6</sub> H <sub>4</sub> S <sub>2</sub> ) <sub>2</sub> | 156,278        | 17,364                                 |                | 17,364           |
| HB(C <sub>6</sub> H <sub>4</sub> S <sub>2</sub> )NH(Et) <sub>2</sub>            |                | 22,985                                 |                | 22,985           |
| (C <sub>6</sub> H <sub>4</sub> )(SH)(SSnBu <sub>3</sub> )                       |                | 276,252                                |                | 27,625           |
| C <sub>6</sub> H <sub>4</sub> (SSnBu <sub>3</sub> ) <sub>2</sub>                |                | 792,217                                |                | 79,222           |
| NH <sub>3</sub> BH <sub>3</sub>   |                |  | 28,354         |                  |
| <b>Total, kg/hr</b>   | <b>340,938</b> | <b>1,489,015</b>                       | <b>196,494</b> | <b>1,295,171</b> |

Reduce mass of reducing agent – reduce cost of recycle

# Can We Get Away Completely From Tin Use?

## Direct Hydrazine Reduction of PB!




# How to Displace Hydrazine and Replace with Ammonia?

## Hydrazine-BH<sub>3</sub> + R → R-BH<sub>3</sub> + Hydrazine Calculations

| Reactant   | B3LYP ΔΔH(rel BDE) |      | G3MP2 ΔΔH(Rel BDE) |      | B3LYP BDE absolute rel to CCSD(T) NH <sub>2</sub> NH <sub>2</sub> |      | G3MP2 BDE absolute rel to CCSD(T) NH <sub>2</sub> NH <sub>2</sub> |      |
|--|--------------------|------|--------------------|------|---|------|---|------|
|  | 0K                 | 298K | 0K                 | 298K | 0K  | 298K | 0K  | 298K |
| Ph-NH <sub>2</sub>                               | 8.0                | 8.2  | 7.0                | 7.3  | 39.0  | 40.4 | 38.0  | 39.5 |
| (C <sub>8</sub> H <sub>18</sub> ) <sub>3</sub> N | 3.0                | 2.9  |                    |      | 34.0  | 35.1 |   |      |
| Et <sub>3</sub> N                                | 3.3                | 3.0  | -1.9               | -2.0 | 34.3  | 35.2 | 29.1  | 30.2 |
| Et <sub>2</sub> NH                               | 0.8                | 0.9  | -1.5               | -1.5 | 31.8  | 33.1 | 29.5  | 30.7 |
| EtMeNH   | -0.7               | -0.6 | -2.6               | -2.5 | 30.4  | 31.6 | 28.4  | 29.7 |
| Me <sub>3</sub> N                                | -0.9               | -0.8 | -4.4               | -4.4 | 30.1  | 31.4 | 26.6  | 27.8 |
| Me <sub>2</sub> NH                               | -1.6               | -1.5 | -3.3               | -3.3 | 29.5  | 30.7 | 27.7  | 28.9 |
| NH <sub>3</sub>                                  | 4.7                | 4.4  | 6.2                | 5.9  | 35.7  | 36.6 | 37.2  | 38.1 |
| Me <sub>2</sub> S                                | 9.4                | 9.8  | 7.8                | 8.3  | 40.4  | 42.0 | 38.8  | 40.5 |
| Et <sub>2</sub> S                                | 8.8                | 9.2  | 6.9                | 7.4  | 39.8  | 41.4 | 37.9  | 39.6 |
| Ph <sub>2</sub> S                                | 14.4               | 14.8 | 12.1               | 12.7 | 45.7  | 47.0 | 43.1  | 44.9 |
| Me <sub>2</sub> O                                | 14.5               | 14.7 | 13.4               | 13.8 | 45.5  | 46.9 | 44.4  | 46.0 |

# Hydrazine-BH<sub>3</sub> + NMe<sub>3</sub>





Pure HzB

Reaction of HzB and Me<sub>3</sub>N in THF overnight:  
~ 60% conversion to Me<sub>3</sub>NBH<sub>3</sub>

Independently synthesized Me<sub>3</sub>NBH<sub>3</sub>  
with excess NH<sub>3</sub> in THF heated at 50°C  
for 5 days: < 5 % conversion to AB




# What About $\text{N}_2\text{H}_4\text{-BH}_3 + \text{NH}_3$ Directly?



Does go, but requires heat:  
at  $60^\circ\text{C}$ : ~ 25% conversion....



# What About PB + Liq. NH<sub>3</sub> in Sealed (One –Pot) System to Drive Reaction?




<sup>11</sup>B-NMR spectrum (in THF) of crude material after venting the NH<sub>3</sub>

i.e. pressure (~ 211 psi) + moderate heat drives the desired rxn.

Can isolate pure samples of AB in > 90% yield using “one–pot” approach.....

# Summary



**"Ideally"**  
 only material consumed would be  $\text{H}_2$   
 (in this case chosen PB = "BNH"  
 (i. e. 2.5 eq.  $\text{H}_2$  released)

**At the inception of CHS CoE (~ 5 yrs ago), many "naysayers" thought that regeneration of AB spent fuel was impossible....**

**Clearly this is not the case.....**

***Remarkable discoveries have been enabled through Center model of collaboration and sharing of information among interdisciplinary experts***

# Acknowledgements

Many DOE Chemical Hydrogen Storage Center of Excellence Collaborators

\$\$\$


DOE EERE



Operated by Los Alamos National Security, LLC for the U.S. Department of Energy's NNSA




# [4+2] Cycloaddition for AB



# How Does This Work? Some Insight From Theory..

**cyclo-B<sub>3</sub>N<sub>3</sub>H<sub>6</sub> (i.e. “BNH<sub>2</sub>”) used as a model fragment i.e. loss of 2 eq. H<sub>2</sub> from AB**  
**Complicated** - many intermediates (only some shown..) on addition(s) of N<sub>2</sub>H<sub>4</sub>



Energies (ΔH<sub>298</sub>) in kcal/mol

DFT/G3MP2B3